




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 257/02	A1	(11) International Publication Number: WO 98/49151 (43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/EP98/02222 (22) International Filing Date: 16 April 1998 (16.04.98) (30) Priority Data: MI97A000982 28 April 1997 (28.04.97) IT (71) Applicant (for all designated States except AU CA GB IE US): BRACCO S.P.A. [IT/IT]; Via E. Folli, 50, I-20134 Milano (IT). (71) Applicant (for AU CA GB IE only): DIBRA S.P.A. [IT/IT]; Piazza Velasca, 5, I-20122 Milano (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): RIPA, Giorgio [IT/IT]; Via E. Folli, 50, I-20134 Milano (IT). ARGESE, Maria [IT/IT]; Via E. Folli, 50, I-20134 Milano (IT). (74) Agents: MINOJA, Fabrizio; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: A PROCESS FOR THE PREPARATION OF 1,4,7,10-TETRAAZACYCLODODECANE <div style="text-align: center;">  </div>		
(57) Abstract A process for the preparation of 1,4,7,10-tetraazacyclododecane of formula (I), starting from decahydro-2a,4a,6a,8a-tetraazacyclopent[fg]acenaphthylene of formula (II), comprising the direct step of hydrolysis in aqueous solution, in conditions of slightly acid, neutral or slightly basic pH, with a primary diamine of formula (V), in which x ranges between 0 and 2, and Q is -CH ₂ CH(OH)CH ₂ -, -(CH ₂) ₂ NH(CH ₂) ₂ -, or -[(CH ₂) ₂ NH] ₂ (CH ₂) ₂ , when x is 1, or Q is -CH ₂ - when x is 2.		


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		


A PROCESS FOR THE PREPARATION OF 1,4,7,10-TETRAAZACYCLO-
DODECANE

The present invention relates to a process for the preparation of 1,4,7,10-tetraazacyclododecane (I) starting from decahydro-2a,4a,6a,8a-tetraazacyclopent-[fg]acenaphthylene (II):



15 The compound of formula (II) can be prepared by cyclization of the intermediate (III), octahydro-3H,6H-2a,5,6,8a-tetraazacenaphthylene, which can in its turn be prepared from triethylenetetramine and glyoxal, as disclosed in International Patent application WO 97/49691 and in Italian Patent application MI96A001257, or from two moles of ethylenediamine through the intermediate of formula (IV), according to the following scheme:


Scheme 1



in which step a) represents the condensation step of triethylenetetramine (or ethylenediamine) with glyoxal, in water or in water-soluble solvents or in mixtures thereof, at a temperature ranging from 0 to 50°C, in the presence of stoichiometric amounts or slight excesses of calcium hydroxide, to give the compound of formula (III) or (IV); and step b) represents, on the other hand, the condensation of the compound of formula (III) or (IV) with an alkylating agent $X-(CH_2)_2-X$, in which X is a halogen or a sulfonic acid reactive derivative, in at least stoichiometric amounts, in the presence of at least 2 moles of a base selected from alkali or alkaline-earth metal carbonates per mole of compound (III), at a temperature of 25-150°C, to give the compound of formula (II).

The intermediate (II) has a remarkable stability in hydrolytic conditions, contrary to the common diaminals which are easily hydrolyzed:


3



For example, publications by Weisman (Tetrahedron Lett., 1980, 21, 335) and by Kolinski (Tetrahedron Lett., 1981, 22, 2217), concerning the synthesis of (II) starting from (I) and glyoxal, confirm the exceptional stability of (II) to acid or basic hydrolytic conditions as well as to reducing agents.

In order to cleave the two carbon atom-bridge which characterizes (II), therefore to obtain (I), an oxidizing process has been described in the already cited Italian Patent application MI96A001257, which allows to transform (II) into oxidation products which can subsequently be transformed into (I) by basic hydrolysis. The whole process from (II) to (I), described in MI96A001257, consists of two steps:

- 1) oxidation of (II)
- 2) hydrolysis of the products from the oxidation of (II)



Alternatively to the oxidative cleavage, WO96/28432

suggests the direct hydrolysis of (II) with hydrobromic acid, or with hydroxylamine in ethanol solution under reflux.

Hydrolysis treatment with hydrobromic acid, not
5 exemplified in WO96/28432, is apparently in disagreement with the teachings of literature, which point out the unexpected stability of compound of formula (II) in acid or basic aqueous solutions.

On the other hand, the reaction with hydroxylamine
10 requires strong excesses of the latter (10 eq. mol) as a free base in ethanol under reflux.


These conditions, although acceptable on laboratory scale, are however not very useful for the industrial processes, as hydroxylamine is quite dangerous
15 (specially when hot) and requires careful handling, use and discharge.

Hydrolysis conditions for (II) which are as selective as possible and, at the same time, the simplest and most cost-saving, should therefore be found
20 to improve the process for the preparation of (I) via (II).

Now it has surprisingly been found, and this is the object of the present invention, a process for the preparation of (I) starting from (II), comprising a
25 hydrolysis step in aqueous solution, in conditions of slightly acid, neutral or slightly basic pH, with a primary diamine of formula (V), as represented in the following Scheme:

5

5



in which

x ranges between 0 and 2, and

- 10 Q is $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2-$, or $-(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2-$, or
 $-[(\text{CH}_2)_2\text{NH}]_2(\text{CH}_2)_2$, when x is 1, or
 Q is $-\text{CH}_2-$ when x is 2.

In the general formula, x can be 0, 1 or 2. In the first case, the diamine corresponds to hydrazine.

- 15 In the other cases, the nature of the diamine depends on the nature of Q, as schematized in the following:

x	Q	Diamine
0	-	Hydrazine
20 1	$-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2-$	1,3-Diamino-2-propanol
1	$-(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2-$	Diethylenetriamine
1	$[(\text{CH}_2)_2\text{NH}]_2(\text{CH}_2)_2$	Triethylenetetramine
2	$-\text{CH}_2-$	Ethylenediamine

- 25 The reaction is extremely selective, allowing not only to obtain high quality (I) in good yields, by also to recycle, in some cases, the reaction by-products, as they can easily be retransformed into (II).

- 30 The reaction takes place in water, in a pH range from 5.5 to 9, preferably from 6 to 8, at temperatures from 60 to 150°C, preferably 60 to 100°C, in the presence of 2-20 mol of diamine per mol of (II), at

6


normal pressure under inert gas atmosphere or in the air, for 12-48 h, or under pressure, for 3-10 h.

At the end of the reaction, the solution is alkalinized with a base, such as sodium hydroxide, concentrated to small volume or to a residue, then (I) is extracted with a suitable solvent, such as toluene, chloroform, butanol, amyl alcohol. The organic phase is concentrated to a residue, to obtain the crude macrocycle (I), which is finally recrystallized from toluene or ethyl acetate.

When the extraction solvent is toluene, it is sufficient to concentrate the toluene solution to a suitable volume, then crystallize the macrocycle (I).

Depending on the diamine used, more developed processes can be defined, providing for the recovery of the reaction by-products.

For example, in the case of triethylenetetramine, the reaction gives (I) and (III), according to the scheme below:




Toluene mother liquors coming from crystallization of (I), substantially containing (III) and triethylenetetramine excess, can be concentrated to a residue, which is in its turn redissolved in water and reacted with glyoxal in the conditions described in

7


MI96A001257, so as to obtain (III), which can optionally be purified, then is transformed into (II) according to the procedure reported in MI96A001257 and finally recycled into the process.

5 In the case of ethylenediamine, the hydrolysis reaction of (II) also gives recyclable by-products:



Compound (IV) can also be recovered from toluene mother liquors and retransformed into (II), as already described in MI96A001257.

15



Some preparation examples according to the method of the invention are reported in the following experimental section

25 **Example 1**

Preparation of (I) by hydrolysis of (II) with diethylenetriamine.

50 g of (II) (0.257 mol) are dissolved in 350 ml of water. 132.5 g of diethylenetriamine (1.285 mol) are added and pH adjusted to 8 with conc. HCl (265 g). The resulting solution is refluxed for 24 h, then cooled,

30

8

alkalinized with NaOH in pellets, concentrated under vacuum to a residual weight of 350 g and extracted (I) with toluene. The combined organic extracts are concentrated to an about 90 mL volume and left to crystallize. 31.5 g of high quality 1,4,7,10-tetraazacyclododecane are obtained (GC titre: 99.5%). Yield. 71%.

Example 2

Hydrolysis of (II) with triethylenetetramine and recovery of the reaction by-products.

50 g of (II) (0.257 mol) are dissolved in 350 ml of water. 105 g (0.640 mol) of triethylenetetramine monohydrate are added and pH is adjusted to 7.5 with conc. HCl. The resulting solution is refluxed for 18 h, then cooled, alkalinized with NaOH in pellets, concentrated under vacuum to a 330 g residual weight and repeatedly extracted with toluene. The combined organic extracts are concentrated under reduced pressure to about 120 mL and left to crystallize. 29.4 g of good quality 1,4,7,10-tetraazacyclododecane are obtained (GC Tit.: >98.5%). Yield: 66%.

Toluene mother liquors are concentrated to a residue, which is redissolved in water. The content in triethylenetetramine and in (III) is determined by GC analysis. Glyoxal and calcium hydroxide are added in amounts such as to transform triethylenetetramine into (III), according to the conditions described in MI 96A 001257. After completion of the reaction, the inorganic salts are filtered off, the filtrate is concentrated to a residue and added with hexane, any insolubles are filtered off and the solution is concentrated to

dryness. 67.2 g (0.40 mol) of (III) are obtained, which can be used to prepare (II) according to the procedure described in MI 96A 001257.

Example 3

5 Hydrolysis of (II) with ethylenediamine and recovery of the reaction by-products.

50 g of (II) (0.257 mol) are dissolved in 300 ml of water. 157.5 g of ethylenediamine (2.57 mol) are added and pH is adjusted to 8 with conc. HCl (310 g).
10 The resulting solution is refluxed for 18 h, then cooled, alkalized with NaOH in pellets, concentrated under reduced pressure to 400 g and repeatedly extracted with toluene. The combined organic extracts are concentrated to a 100 mL volume and left to crystallize.
15 24.8 g of high quality 1,4,7,10-tetraazacyclododecane are obtained (GC titre: >99%). Yield: 56%.

Toluene mother liquors are concentrated under reduced pressure. The residue is subjected to chromatography on silica (eluent: chloroform: methanol:25% ammonia = 6:3:1). 21 g (0.148 mol) of (IV)
20 are obtained, which can be retransformed into (II) in the conditions described in MI 96A 001257.

Example 4


25 Hydrolysis of (II) with various diamines at various pH and temperatures.

19.4 g (0.1 mol) of (II) are hydrolyzed with a similar procedure to that described in example 1, using the diamines and the conditions reported in the following table:

	Diamine	mol/mol (II)	10		t (h)	% Yield
			pH	T (°C)		
5	Hydrazine	20	7	100	24	60%
	Hydrazine	10	6.5	100	48	55%
	Ethylene- diamine	10	6.5	100	18	52%
	Ethylene- diamine	8	7	80	24	45%
10	Ethylene- diamine	10	5.5	100	18	42%
	Diethylene- triamine	4	7.5	80	24	58%
	Diethylene- triamine	6	6.5	100	12	62%
	Diethylene- triamine	6	8	60	48	65%
15	Triethylene- tetramine	4	6	80	24	55%
	Triethylene- tetramine	2	8	100	48	62%
	Triethylene- tetramine	2	9	100	48	57%
	1,3-Diamino- 2-propanol	6	7	100	24	51%
20	1,3-Diamino- 2-propanol	5	8	100	18	48%

CLAIMS

1. A process for the preparation of 1,4,7,10-tetraazacyclododecane of formula (I) starting from
 5 decahydro-2a,4a,6a,8a-tetraazacyclopent[fg]acenaphthylene of formula (II):



- 15 comprising a direct step of hydrolysis in aqueous solution, in conditions of slightly acid, neutral or slightly basic pH, with a primary diamine of formula (V),

wherein:

- x ranges between 0 and 2, and
 20 Q is $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2-$, or $-(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2-$, or
 $-\text{[(CH}_2)_2\text{NH}]_2(\text{CH}_2)_2$, when x is 1, or
 Q is $-\text{CH}_2-$ when x is 2.

2. A process according to claim 1, in which the
 25 diamine is selected from the group of: diethylenetriamine, triethylenetetramine and ethylenediamine.


3. A process according to claims 1-2, in which the
 hydrolysis reaction is carried out in water, at a pH
 from 5.5 to 9, at temperatures ranging from 60 to 150°C,
 preferably 60 to 100°C, in the presence of 2-20 mol
 30 of diamine per mole of (II), at normal pressure under
 inert gas atmosphere or in the air, for 12-48 h,

12

or under pressure, for 3-10 h.

4. A process according to claim 3, in which the reaction pH ranges from 6 to 8.

5. A process according to claims 1-4, in which the
 5 diamine is triethylenetetraamine, represented in the following scheme




in which compound of formula (III), octahydro-3H,6H-
 2a,5,6,8a-tetraazacenaphthylene, is simultaneously
 15 formed, which is an intermediate useful for the
 preparation of the compound of formula (II), decahydro-
 2a,4a,6a,8a-tetraazacyclopent[fg]acenaphthylene,
 according to known methods.

6. A process according to claim 5, in which the mother
 20 liquors from the crystallization of the compound of
 formula (I), substantially containing the compound of
 formula (III) and the triethylenetetraamine excess, are
 concentrated to a residue, which is in its turn
 redissolved in water and can be used as an intermediate
 25 for the preparation of the compound of formula (II),
 according to known methods.

7. A process according to claims 1-4, in which the
 diamine is diethylenetriamine.

8. A process according to claims 1-4, in which the
 30 diamine is ethylenediamine, according to the following
 scheme

13



in which compound of formula (IV), decahydro-pyrazino[2,3-b]pyrazine, is simultaneously formed, which is an intermediate useful for the preparation of the compound of formula (II), decahydro-2a,4a,6a,8a-tetra-azacyclopent[fg]acenaphthylene, according to known methods.

9. A process according to claim 8, in which the mother liquors from the crystallization of compound of formula (I), substantially containing compound of formula (IV), are concentrated to a residue, which is in its turn redissolved in water and can be used as an intermediate for the preparation of compound of formula (II), according to known methods.

20

INTERNATIONAL SEARCH REPORT

national Application No
PCT/EP 98/02222

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D257/02

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 28432 A (NYCOMED IMAGING A/S) 19 September 1996 cited in the application see claims ----	1-9
A	EP 0 427 595 A (L'AIR LIQUIDE) 15 May 1991 see claims ----	1-9
P,A	WO 97 49691 A (BRACCO S.P.A. ET AL.) 31 December 1997 cited in the application see claims -----	1-9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 August 1998

Date of mailing of the international search report

26/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/02222

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9628432	A	19-09-1996	AU 4838996	A	02-10-1996
			CA 2214991	A	19-09-1996
			CN 1183776	A	03-06-1998
			EP 0815090	A	07-01-1998
			NO 974169	A	07-11-1997
			US 5589595	A	31-12-1996
<hr/>					
EP 427595	A	15-05-1991	FR 2654102	A	10-05-1991
			AU 649632	B	02-06-1994
			AU 6592790	A	25-07-1991
			CA 2029454	A	10-05-1991
			JP 3169870	A	23-07-1991
			US 5434262	A	18-07-1995
<hr/>					
WO 9749691	A	31-12-1997	IT MI961257	A	22-12-1997
			AU 3256497	A	14-01-1998
<hr/>					